

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 209/88, A61K 31/40	A1	(11) International Publication Number: WO 99/05105 (43) International Publication Date: 4 February 1999 (04.02.99)
(21) International Application Number: PCT/EP98/04475 (22) International Filing Date: 18 July 1998 (18.07.98) (30) Priority Data: 97112491.2 22 July 1997 (22.07.97) EP (71) Applicant (for all designated States except US): BOEHRINGER MANNHEIM GMBH [DE/DE]; D-68298 Mannheim (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): REINHOLZ, Erhard [DE/DE]; Verl. Triebstrasse 1, D-68542 Heddesheim (DE). BEYER, Peter [DE/DE]; Im Kirchenstück 8, D-67433 Neustadt (DE). (74) Common Representative: BOEHRINGER MANNHEIM GMBH; Patentabteilung, D-68298 Mannheim (DE).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: THERMODYNAMICALLY STABLE MODIFICATION OF 1-(4-CARBAZOLY-LOXY)-3-[2-(2-METHOXYPHENOXY)ETHYLAMINO]-2-PROPANOLE, PROCESS FOR ITS PREPARATION AND PHARMACEUTICAL COMPOSITIONS CONTAINING IT		
(57) Abstract <p>The present invention relates to a new thermodynamically stable modification of 1-(4-carbazolyloxy)-3-[2-(2-methoxyphenoxy)ethylamino]-2-propanole (Carvedilol), pharmacologically acceptable salts, or optically active forms thereof, processes for the preparation, and pharmaceutical compositions containing it.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5 **THERMODYNAMICALLY STABLE MODIFICATION OF**
1-(4-CARBAZOLYLOXY)-3-[2-(2-METHOXYPHENOXY)ETHYLAMINO]-
2-PROPANOLE, PROCESS FOR ITS PREPARATION AND
PHARMACEUTICAL COMPOSITIONS CONTAINING IT

10

Description

The present invention relates to a new thermodynamically stable modification of (\pm)1-(4-carbazolyloxy)-3-[2-(2-methoxyphenoxy)ethylamino]-2-propanole (Carvedilol), pharmaco-
15 logically acceptable salts or optically active forms thereof, processes for the preparation, and pharmaceutical compositions containing it.

Carvedilol, having a melting point of 114-115°C, is a compound with excellent pharmacological properties (Merck Index 11. Ed. No. 1882), known to be active in the treatment of cardiac diseases. The preparation and its use in medicine is described in EP-B-0
20 004 920.

Carvedilol has a chiral center and, as such, can exist either as individual stereoisomers or in racemic form. Both the racemate and stereoisomers may be obtained according to
25 procedures well known in the art (EP-B-0127099).

It has now been discovered that Carvedilol can be isolated in two different modifications depending upon the method of preparation which are distinguishable by their infra-red Raman and X-ray powder diffraction spectra, and their melting points. The two polymorphic forms are monotropic and they are hereinafter designated as Form I and Form
30

II. It is desirable to prepare a therapeutic agent consisting of an unique and defined composition which has a high storage stability.

The present invention provides a thermodynamically stable crystalline form of Carvedilol
5 substantially free of other physical forms having a melting point about 123-126°C, and an
infra-red spectrum with a sharp peak at 3451 cm⁻¹, which is referred to hereinafter as
Form I.

The invention also provides a process for producing this substantially pure Form I. In
10 another embodiment of this invention, there is provided a pharmaceutical formulation
containing the substantially pure Form I of Carvedilol as an active ingredient.

Finally, the present invention provides a method of using the new substantially pure form
to prevent and/or treat circulatory and cardiac diseases.
15

Where reference is made in this application to Form I or Form II substantially free of
other physical forms, it preferably means that at least 90% by weight of Form I or Form
II is present in that modification.

20 Form II is the modification of Carvedilol prepared and purified according to EP-B-0 004
920.

Surprisingly it was now found that a new thermodynamically stable modification of
Carvedilol (Form I) with a higher melting point is obtained when the process of manu-
25 facture is slightly altered.

The melting points of each Forms I respectively II depend upon their level of purity,
consequently Form I has been found to have a melting point of about 123-126°C, Form
II about 114-115°C.
30

Furtheron it has been discovered that Form I is that of being the thermodynamically stable form, which is of advantage. Therefore this thermodynamically stable form is given preference in the preparation of pharmaceutical formulations.

5 Pharmaceutically acceptable salts are considered to be encompassed within the compounds and the method of the present invention. The term „pharmaceutically acceptable salts“ refers to salts of substantially pure Form I which are substantially non-toxic to living organism. For the conversion of Carvedilol into its pharmacologically acceptable salts, it is reacted, preferably in an organic solvent, with an equivalent amount of an
10 inorganic or organic acid, for example hydrochloric acid, hydrobromic acid, phosphoric acid, sulphuric acid, acetic acid, citric acid, maleic acid or benzoic acid. It should be recognized that any particular anion forming a part of any salt of this invention is not critical, so long as the salt, as a whole, is pharmacologically acceptable and as long as the anion moiety does not contribute undesired qualities.

15

For the resolution of the racemates, there can be used for example, tartaric acid, malic acid, camphoric acid or camphorsulphonic acid.

According to another aspect, the invention provides a pharmaceutical composition,
20 which comprises Form I substantially free of other physical forms and a pharmaceutical acceptable carrier or adjuvant.

The compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route and in
25 dose effective for the treatment intended. Therapeutically effective doses of the compounds of the present invention required to prevent or arrest the progress of the medical condition are readily ascertained by one of ordinary skill in the art.

Accordingly, the invention provides a class of novel pharmaceutical compositions
30 comprising Carvedilol of Form I of the present invention, in association with one or more non-toxic pharmaceutically acceptable carriers and/or adjuvants (collectively referred to

herein as „carrier materials“) and, if desired, other active ingredients. The compounds and compositions may, for example, be administered orally, intravascularly, intraperitoneally, subcutaneously, intramuscularly or topically.

- 5 For all administrations, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, creme, ointment, gel, lotion, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. A suitable daily dose for a mammal may vary widely depending on the condi-
- 10 tion of the patient and other factors. However, a dose from about 0.01 to 100 mg/kg body weight, particularly from about 0.05 to 3 mg/kg body weight, respectively 0.01-10 mg/cm² skin, may be appropriate. The active ingredient may also be administered by injection.
- 15 The dose regimen for treating a disease condition with the compounds and/or compositions of this invention is selected in accordance with a variety of factors, including the type, age, weight, sex and medical conditions of the patient and in accordance to the severity of the disease and thus may vary widely.
- 20 For therapeutic purposes, the compounds of the invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If per os, the compound may be admixed with lactose, sucrose, starch powder, cellulose esters of alcanoic acids, cellulose alkyl ester, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulphuric acids, gelatine, acacia,
- 25 sodium alginate, polyvinyl-pyrrolidone and/or polyvinyl alcohol, and thus tableted or encapsulated for convenient administration. Alternatively, the compound may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cotton seed oil, peanut oil, sesam oil, benzyl alcohol, sodium chloride and/or various buffers. Appropriate additives for the use as ointments, cremes or gels are for example paraffine,
- 30 vaseline, natural waxes, starch, cellulose, or polyethylenglycole. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

Appropriate dosages in any given instance, of course, depend upon the nature and severity of the condition treated, the route of administration and the species of mammal involved, including its size and any individual idiosyncracies.

- 5 Representative carriers, dilutions and adjuvants include, for example, water, lactose, gelatine starch, magnesium stearate, talc, vegetable oils, gums, polyalkylene glycols, petroleum jelly, etc. The pharmaceutical compositions may be made up in a solid form, such as granules, powders or suppositories, or in liquid form, such as solutions, suspen-
10 sions or emulsions. The pharmaceutical compositions may be subjected to conventional pharmaceutical adjuvants, such as preservatives, stabilizers, wetting agents, emulsifiers, buffers, etc.

As indicated, the dose administered and the treatment regimen will be dependent, for example, on the disease, the severity thereof, on the patient being treated and his
15 response to treatment and, therefore, may be widely varied.

Characterization of Forms I and II of Carvedilol

20 Thermomicroscopy

Thermal analysis was carried out with a Kofler heating stage (Reichert, Vienna) mounted on a video-equipped Olympus microscope BH-2 or with a Kofler heating stage microscope Thermovat[®] (Reichert, Vienna); both microscopes with polarisation facility and
25 digital thermometer.

Form II consists of heterogeneously looking rhombohedral to hexagonally shaped lamellar crystals up to a size of 120 μm , which melt about 114-115°C, whereas Form I consists of 40 μm large grains, respectively prisms, which melt about 123-126 °C.

30

Differential Scanning Calorimetry (DSC)

DSC was carried out with DSC-7 (Perkin-Elmer, Norwalk, Ct., USA) equipped with cooling system CCA-7, perforated Al sample capsules (25 μ l), weighed object 1.5 mg each (ultra-micro weighing scale UM 3, Mettler, CH-Greifensee, Switzerland). Nitrogen 4.0 as flushing gas (20 ml min⁻¹). Computer-aided recording of DSC signal. Calibration of temperature indication for CCA curves with melting point for water and caffeine anhydrate (melting point 236.2 °C), each with tightly sealed sample capsule. Calibration of ordinates (DSC signal) with melting heat of indium 99.999% (Perkin-Elmer, Norwalk, Ct., USA).

10

The measured melting points correspond to the ones determined thermomicroscopically. It could be estimated by the way of the measured melting heats (Form I: ΔH_f 48.2 kJ/mol; Form II ΔH_f 51.0 kJ/mol), that the crystallisate consisting of Form I is contaminated with approximately 2 to 3% of Form II, which could also be seen thermomicroscopically.

15

FT-IR, FT Raman spectroscopy and X-ray diffractometry

FT-IR spectroscopy was carried out with a Bruker IFS 25 FT-IR spectrometer. For the production of the KBr compacts approximately 1 mg of sample was powdered with 270 mg of KBr. The spectra were recorded in transmission mode ranging from 4000 to 600 cm⁻¹. Resolution: 2 cm⁻¹ (50 interferograms).

20

FT Raman spectroscopy was carried out with Bruker RFS 100 FT Raman spectrometer, equipped with a diode-pumped Nd:YAG laser (1064 nm) and a liquid nitrogen cooled highly sensitive detector. The powdered samples were pressed into small aluminium fittings, the spectra were recorded at an initial capacity of 200 mW, resolution: 4 cm⁻¹ (50 interferograms).

25

X-ray powder diffractometry was carried out with a Simens X-ray diffractometer D-5000, Diffrac/AT with θ/θ goniometer, CuK α -rays, nickel filter for monochromatisation,

30

rotation of sample during measurement, scintillation counter, angular range 2° to 40° (2θ), steps of 0.01° (2θ), measuring time 2 secs.

The IR spectra of both modifications show great differences in the stretching vibration range (Form I 3451 cm^{-1} ; Form II 3345 cm^{-1})(Fig. 1,2), which are caused by different hydrogen bridges. This corresponds to the Raman spectra differing only little. The biggest difference in the Raman spectra is at approximately 2942 and approximately 755 cm^{-1} (Fig.3,4). The X-ray powder diffraction pattern of Form I has characteristic peaks occurring at $2\theta = 9.5, 10.8, 12.0, 14.6, 19.6, 21.5,$ and 22.3 (Fig. 5) whereas the characteristic peaks of Form II occur at $2\theta = 5.9, 14.9, 17.6, 18.5,$ and 24.4 (Fig. 6).

Process for preparing Form I Carvedilol

Example 1

15

Crude Carvedilol is prepared according to the procedure described in EP-B-0 004 920, in methanol. Crude Carvedilol (based on 300 g dry Carvedilol), 15 g CXA-coal and 2800 ml methanol are heated for 15 minutes under reflux in a three-neck-flask. The hot solution is filtered and washed with 300 ml hot methanol and heated under reflux again. Subsequently the solution is cooled down during half an hour to 30°C , stirred between 3 to 22 hours and cooled down slowly to 0°C in $3\frac{1}{2}$ hours. After stirring the solution for additional two hours at 0°C the product is isolated, washed three times with 40 ml methanol and dried under vacuo at 60°C for 24 hours. 203-255 g of pure Form I are obtained and characterized as described before.

25 Form II can be obtained by an additional recrystallization process in isopropanol.

Example 2

A 1:1 mixture of Form I and Form II was suspended in isopropanol and agitated with a magnetic stirrer for 18 h in a tightly sealed glass cylinder. During this time the temperature was repeatedly increased and lowered between 10 and 25°C . Subsequently the


sample was filtered with a micro glass filter funnel (G3), dried and evacuated. The IR spectrum of this sample corresponds to Form I. The DSC curve does not show a peak between 114-115°C, thus this is pure Form I.

- 5 It will be understood that the specification and examples are illustrative but not limitative of the present invention and that other embodiments within the spirit and scope of the invention will suggest themselves to those skilled in the art.


Claims

1. Modification of (\pm)-1-(4-carbazolyloxy)-3-[2-(2-methoxyphenoxy)ethylamino]-2-propanole having the following X-ray diffraction pattern obtained with a $\text{CuK}\alpha$ radiation at $2\theta = 9.5, 10.8, 12.0, 14.5, 19.6, 21.5, 22.3$, and an infrared spectrum having sharp peaks at 3451 cm^{-1} , wherein the melting point is about $123\text{--}126^\circ\text{C}$.
5
2. Pharmacologically acceptable salts or optical active forms of the compound according to claim 1.
10
3. A process for preparing and isolating substantially pure form of the compound as claimed in claim 1.
15
4. The process of claim 3, wherein the rearrangement is carried out at a temperature between 25 and 35°C for a time from 3 to 22 hours in methanol.
5. The process of claim 3 or 4, wherein the higher melting carvedilol modification is recovered at 0°C from the rearrangement reaction mixture.
- 20 6. A pharmaceutical composition comprising a substantially pure form of the stable modification of 1-(4-carbazolyloxy)-3-[2-(2-methoxyphenoxy)ethylamino]-2-propanole as the active ingredient as claimed in claim 1 together with one or more pharmaceutical acceptable carriers or adjuvants.
- 25 7. The use of a pharmaceutical composition as claimed in claim 5 for the manufacturing of a medicament for the prophylaxis or treatment of cardiac diseases.


Figur 1.




Figur 2




Figur 3




Figur 4



Figur 5



Figur 6



INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/04475

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D209/88 A61K31/40

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 127 099 A (BOEHRINGER MANNHEIM) 5 December 1984 cited in the application see claims; examples 7,8 ---	1-7
A	DE 28 15 926 A (BOEHRINGER MANNHEIM) 31 October 1979 cited in the application see claims; example 2 ---	1-7
A	G. Z. FEUERSTEIN ET. AL.: "carvedilol update III. Rationale for use in congestive heart failure." DRUGS TODAY, vol. 31F, September 1995, pages 1-23, XP002039368 see whole document -----	1-7

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

15 October 1998

Date of mailing of the international search report

02/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Helps, I

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 98/04475

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0127099 A	05-12-1984	DE 3319027 A	29-11-1984
		AU 551116 B	17-04-1986
		AU 2848084 A	29-11-1984
		CA 1259071 A	05-09-1989
		CA 1257279 C	11-07-1989
		DK 91393 A	06-08-1993
		DK 255184 A	27-11-1984
		FI 842046 A,B,	27-11-1984
		IE 57533 B	24-03-1993
		JP 1818634 C	27-01-1994
		JP 5027622 B	21-04-1993
		JP 59222473 A	14-12-1984
		JP 1917129 C	23-03-1995
		JP 5208957 A	20-08-1993
		JP 6013508 B	23-02-1994
		PT 78633 B	18-06-1986
		US 4824963 A	25-04-1989
		US 4985454 A	15-01-1991
		US 4697022 A	29-09-1987
		US 5071868 A	10-12-1991
DE 2815926 A	18-10-1979	AT 375639 B	27-08-1984
		AU 522975 B	08-07-1982
		AU 4582079 A	18-10-1979
		BG 61419 B	31-07-1997
		CA 1129416 A	10-08-1982
		CS 227007 B	16-04-1984
		CS 9104200 A	15-04-1992
		CS 227047 B	16-04-1984
		DD 143607 A	03-09-1980
		DK 141979 A,B,	14-10-1979
		EP 0004920 A	31-10-1979
		FI 791142 A,B,	14-10-1979
		HK 2385 A	18-01-1985
		JP 1023462 B	02-05-1989
		JP 1545837 C	28-02-1990
		JP 54157558 A	12-12-1979
		JP 63258416 A	25-10-1988
		LT 112493 R	25-04-1994
		LU 88320 A	04-05-1994

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/04475

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2815926 A		MX 9203380 A	01-09-1992
		US 4503067 A	05-03-1985
		ZA 7901732 A	28-05-1980
<hr/>			